



Complete Summary

GUIDELINE TITLE

Treatment of obesity.

BIBLIOGRAPHIC SOURCE(S)

Finnish Medical Society Duodecim. Treatment of obesity. In: EBM Guidelines. Evidence-Based Medicine [CD-ROM]. Helsinki, Finland: Duodecim Medical Publications Ltd.; 2004 Jun 22 [Various].

GUIDELINE STATUS

This is the current release of the guideline.

COMPLETE SUMMARY CONTENT

SCOPE

METHODOLOGY - including Rating Scheme and Cost Analysis

RECOMMENDATIONS

EVIDENCE SUPPORTING THE RECOMMENDATIONS

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

IMPLEMENTATION OF THE GUIDELINE

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT

CATEGORIES

IDENTIFYING INFORMATION AND AVAILABILITY

DISCLAIMER

SCOPE

DISEASE/CONDITION(S)

Obesity

GUIDELINE CATEGORY

Counseling

Evaluation

Management

Treatment

CLINICAL SPECIALTY

Family Practice

Internal Medicine

Nutrition

INTENDED USERS

Dietitians
Health Care Providers
Nurses
Physicians

GUIDELINE OBJECTIVE(S)

Evidence-Based Medicine Guidelines collect, summarize, and update the core clinical knowledge essential in general practice. The guidelines also describe the scientific evidence underlying the given recommendations.

TARGET POPULATION

Patients who are obese

INTERVENTIONS AND PRACTICES CONSIDERED

Evaluation

1. Assess the need for treatment using body mass index (BMI)

Counseling/Treatment

1. Basic treatment (lifestyle counseling regarding behaviour, diet, and exercise)
2. Basic treatment plus a very low calorie diet
3. Drug (orlistat or sibutramine)
4. Surgical treatment (gastroplasty, gastric banding, or gastric bypass)
5. Follow-up

Note: Guideline developers considered but did not specifically recommend guar gum, ephedrine, ephedrine plus caffeine, dietary supplements containing ephedra, or chitosan.

MAJOR OUTCOMES CONSIDERED

- Morbidity associated with obesity
- Efficacy of treatment
 - Weight loss
 - Maintenance of weight loss
 - Amount of weight regained
 - Prevention of progression of obesity
 - Effect of treatment on comorbidities
- Adverse effects or complications associated with treatment

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Hand-searches of Published Literature (Primary Sources)
Hand-searches of Published Literature (Secondary Sources)
Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

The evidence reviewed was collected from the Cochrane database of systematic reviews and the Database of Abstracts of Reviews of Effectiveness (DARE). In addition, the Cochrane Library and medical journals were searched specifically for original publications.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Weighting According to a Rating Scheme (Scheme Given)

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Levels of Evidence

- A. Strong research-based evidence. Multiple relevant, high-quality scientific studies with homogenic results.
- B. Moderate research-based evidence. At least one relevant, high-quality study or multiple adequate studies.
- C. Limited research-based evidence. At least one adequate scientific study.
- D. No research-based evidence. Expert panel evaluation of other information.

METHODS USED TO ANALYZE THE EVIDENCE

Review of Published Meta-Analyses
Systematic Review

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Not stated

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

COST ANALYSIS

Cost analyses were reviewed.

In a cost-utility analysis, the net cost per quality-adjusted life years (QALY) was 6,289 Great Britain pound GBP for gastric bypass (Roux-en-Y), 10,237 GBP for vertical banded gastroplasty, and 8,527 GBP for silicone adjustable gastric banding as compared with non-surgical management. Gastric bypass appears to have a modest net cost per QALY gained compared to vertical banded gastroplasty (742 GBP/QALY), when silicone adjustable gastric banding has a large net cost per QALY gained compared to gastric bypass (256,856 GBP/QALY).

METHOD OF GUIDELINE VALIDATION

Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Not stated

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

The levels of evidence [A-D] supporting the recommendations are defined at the end of the "Major Recommendations" field.

References: (Karvetti & Hakala, 1992; Wadden, 1993; Garrow, 1992; Anderson, Hamilton, & Brinkman-Kaplan, 1992)

Assessment of the Need of Treatment

- Overweight (mild obesity): body mass index (BMI) 25 to 30 kilograms (kg)/m²
 - Usually no treatment is indicated.
 - Treatment (Glenny & O'Meara, 1997; DARE-971098, 1999) [A] is indicated in central obesity, metabolic syndrome (see Finnish Medical Society Duodecim guideline "Metabolic Syndrome [MBS]"), or non-insulin-dependent diabetes.
 - In children even mild obesity should be treated.
- Moderate obesity: BMI 30-35 kg/m²
 - Treatment (Glenny & O'Meara, 1997; DARE-971098, 1999) [A] is always indicated if the patient has diabetes, hypertension, hyperlipidaemia, or other associated disease (Douketis et al., 1999; DARE-998429, 2000) [B].
 - A young obese person with good health should be treated. The treatment of middle-aged persons is decided individually according to the available resources.
- Severe obesity: BMI >35 kg/m²
 - Must always be treated

Selecting the Method of Treatment

1. Basic treatment consisting of a gradual and permanent change in living habits by counselling and guidance.
 - Well suited for patients with mild and moderate obesity and for the majority of patients with severe obesity; should always be included in other forms of conservative treatment
2. Basic treatment and a very low calorie diet
 - Well suited for morbid and severe obesity
 - A choice in moderate obesity if the basic treatment has been unsuccessful and there is a strong indication for reducing weight (associated diseases)
3. Drugs (orlistat or sibutramine)
 - Do not automatically help all patients
 - An alternative especially if other approaches have been failed
 - Life-style counselling must also be included.
4. Surgical treatment
 - Suitable only for selected patients with morbid obesity (see criteria below).

Basic Treatment

Organisation

- Group treatment is less costly and as effective as individualised treatment
 - At least 10 meetings are arranged with about one-week intervals.
 - The group leader is a nurse or a dietician with special training in the treatment of obesity.

Goals that Can Be Measured

- The optimal rate of weight reduction is 0.5 kg/week. As the adipose tissue contains about 30 MJ (7,000 kcal)/kg, a daily reduction of 2,100 kJ (500 kcal) in the energy intake will result in this rate of weight reduction.
- The goal is to reduce weight by 5 to 10%, which already results in significant benefit in the treatment of diseases associated with obesity.
- A permanent result is always aimed at. This means that the changes in living habits must be permanent.
- In growing children, weight should be kept constant so that the growth of height corrects the relative weight.
- There are many treatments with no proven efficacy. Appetite-suppressant drugs may result in a moderate weight reduction but the effect is transient.

Aims and Contents of Counselling

- Changes in knowledge and attitudes
 - Human energy consumption is reduced if the body weight is reduced. In order to sustain the weight that has been achieved the changes in living habits must be permanent.
- Changes in meals (Miller, Koceja, & Hamilton, 1997; DARE-971214, 1999) [A]
 - Find out the present contents of meals.
 - Reduce intake by about 2,000 kJ (500 kcal) daily.

- The main emphasis is on the reduction of fat intake (Pirozzo et al., 2002) [B].
- Remember alcohol as a cause of obesity.
- Small daily changes are effective in the long run.
- Keep three daily meals.
- Changes in physical exercise (Miller, Koceja, & Hamilton, 1997; DARE-971214) [A]
 - The advice depends on the degree of obesity.
 - Exercise during daily activities should be encouraged (climbing up stairs, walking or cycling to work).
- Changes in eating behaviour
 - The most common goal is to change behaviour, not to "hunt kilograms."
 - Identify the circumstances that trigger eating.
 - Grocery shopping with a pre-planned list
 - Reduce temptations (no food in sight!).
 - Do nothing else when eating (such as watch TV, read magazines).
 - Eat slowly.

Very Low Calorie Diet (VLCD)

- Reference: (Anderson, Hamilton, & Brinkman-Kaplan, 1992)
- Constituents
 - 1,700 to 2,100 kJ (400 to 500 kcal) of energy, a maximum of 3,300 kJ (800 kcal) daily
 - Protein as needed (at least 50 grams (g) daily)
 - Essential fatty acids, trace elements, and vitamins as needed
- Schedule
 - Ready-made commercial formulas should be used as the only food continuously for 8 to 10 weeks in severe obesity and for a shorter period in milder obesity, but usually for a minimum of 6 weeks.
 - The patient is followed up at 1 to 2 week intervals.
 - Suitable for patients with non-insulin-dependent (type 2) diabetes and hypertension. Insulin treatment is stopped or the dose is reduced considerably, and the dose of sulphonylureas is halved (risk of hypoglycaemia!) before starting very low calorie diet. The doses of other drugs need not be reduced.
 - The rate of weight reduction is about 1.5 to 2 kg/week, and the short-term weight reduction is 2 to 2.5 times that on basic treatment.
 - A very low calorie diet alone does not yield permanent results. Basic treatment for permanent life-style changes is applied in the normal fashion.

Weight-Reducing Drugs

- Orlistat is a lipase inhibitor. The most common side effects are bloating and diarrhoea after fatty meals.
- Sibutramine is a centrally acting appetite suppressant. Side effects are slight increase of blood pressure and pulse rate, dry mouth, and constipation.
- With both drugs mean weight reduction is 4 to 5 kg better than with placebo (Padwal, Li & Lau, 2003) [B].

- Stopping of a drug leads to relapse (i.e., permanent result needs continuous use).
- Do not use alone. Life-style counselling is always needed, and, with orlistat, reduction of food fat content is mandatory.
- Drugs are not a firsthand treatment of obesity. They can be used if other approaches have failed.
- If no significant weight reduction can be seen over 2 to 3 months, stop the drug.

Surgical Treatment

Criteria

- Age below 60 years
- BMI at least 35 to 40 kg/m².
- An efficient conservative treatment strategy has been tried.
- The patient is cooperative.
- There is no abuse of alcohol or drugs.

Method

- Gastroplasty, gastric banding (Schneider, 2000; DARE-20018109, 2002; Chapman et al., 2002; HTA-20030005, 2004; "Newer techniques," 2003; HTA-20031130, 2004) [B], or gastric bypass so that the patient can eat only slowly and small amounts at a time. There are several surgical techniques, including laparoscopic procedures.
- The operation is not sufficient alone. Adequate preoperative investigations, patient counselling, and organized follow-up are mandatory.
- The outcome of successful surgical treatment is much better than that of conservative treatment (Colquitt et al., 2003; Clegg et al., 2002; HTA-20020727, 2004; "Special report," 2003; HTA-20031129, 2004; "Gastric restrictive surgery," 2000; HTA-20030550, 2004; "Newer techniques," 2003; HTA-20031130, 2004; Nilsen, 2003; HTA-20030502, 2004) [A]: the patients reduce 30 to 40 kg of their weight, and the result is long lasting (Glenny & O'Meara, 1997; DARE-971098, 1999) [A].
- Some patients experience complications after surgery.

Related Evidence

- Guar gum is not effective in weight reduction and causes adverse effects (Pittler & Ernst, 2001; DARE-20011626, 2002) [A].
- Ephedrine, ephedrine plus caffeine, or dietary supplements containing ephedra may be effective in producing a modest short-term weight loss as compared to placebo (Agency for Healthcare Research and Quality, 2003; HTA-20030424, 2004) [C].
- Chitosan may be more effective than placebo in the short-term treatment of overweight and obesity, but the evidence comes mainly from poor quality studies (Ni Mhurchu, 2005)

Definitions:

Levels of Evidence

- A. Strong research-based evidence. Multiple relevant, high-quality scientific studies with homogenic results.
- B. Moderate research-based evidence. At least one relevant, high-quality study or multiple adequate studies.
- C. Limited research-based evidence. At least one adequate scientific study.
- D. No research-based evidence. Expert panel evaluation of other information.

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

REFERENCES SUPPORTING THE RECOMMENDATIONS

[References open in a new window](#)

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

Concise summaries of scientific evidence attached to the individual guidelines are the unique feature of the Evidence-Based Medicine Guidelines. The evidence summaries allow the clinician to judge how well-founded the treatment recommendations are. The type of supporting evidence is identified and graded for select recommendations (see the "Major Recommendations" field).

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

Appropriate assessment and treatment of obesity

POTENTIAL HARMS

- The most common side effects of orlistat are bloating and diarrhoea after fatty meals.
- Side effects of sibutramine are slight increase of blood pressure and pulse rate, dry mouth, and constipation.
- The complications reported from laparoscopic adjustable gastric banding included aspiration pneumonia, band slippage, and rotated or infected access ports. Band migration was more common if the band injections were performed by residents or nurses, rather than senior surgeons.
- Some complications of gastric surgery can occur, such as wound infection.

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Getting Better
Living with Illness

IOM DOMAIN

Effectiveness
Patient-centeredness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Finnish Medical Society Duodecim. Treatment of obesity. In: EBM Guidelines. Evidence-Based Medicine [CD-ROM]. Helsinki, Finland: Duodecim Medical Publications Ltd.; 2004 Jun 22 [Various].

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

2004 Jun 22

GUIDELINE DEVELOPER(S)

Finnish Medical Society Duodecim - Professional Association

SOURCE(S) OF FUNDING

Finnish Medical Society Duodecim

GUIDELINE COMMITTEE

Editorial Team of EBM Guidelines

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Primary Author: Pertti Mustajoki

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

Not stated

GUIDELINE STATUS

This is the current release of the guideline.

GUIDELINE AVAILABILITY

This guideline is included in a CD-ROM titled "EBM Guidelines. Evidence-Based Medicine" available from Duodecim Medical Publications, Ltd, PO Box 713, 00101 Helsinki, Finland; e-mail: info@ebm-guidelines.com; Web site: www.ebm-guidelines.com.

AVAILABILITY OF COMPANION DOCUMENTS

None available

PATIENT RESOURCES

None available

NGC STATUS

This NGC summary was completed by ECRI on September 2, 2005.

COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers

or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2006 National Guideline Clearinghouse

Date Modified: 9/25/2006

